

Enzyme Potentiated Hyposensitization

L. M. McEWEN, B.M., B. CH.

Effect of glucose, glucosamine, N-acetylamino-sugars and gelatin on the ability of F Beta glucuronidase to block the anamnestic response to antigen in mice.

Abstract: In mice sensitized to horse serum, Beta glucuronidase prevents the increased sensitivity which results from a second dose of antigen given one week before challenge by pinnal anaphylaxis. Beta glucuronidase loses this activity with age or in the presence of gelatin. In both cases glucose will restore the immunological blocking activity of the enzyme. These results suggest a reason for variability in the hyposensitizing activity of the different samples of enzyme used in earlier clinical and animal experiments.

Introduction

Following the discovery of the ability of the enzyme Beta glucuronidase to potentiate the hyposensitizing effect of small doses of antigen in the treatment of clinical allergy the first paper in this series described a similar behaviour of the enzyme in three different models of experimental anaphylaxis.

However, the potentiating effect of different samples of Beta glucuronidase has been found to vary, so that it was impractical to proceed to further clinical trials and the results of animal experiments were not consistent.

This paper is a study of the source of the variability and describes the effect of glucose, glucosamine, certain N-acetyla-

mino-sugars and gelatin on the ability of the enzyme to modify the anamnestic response to a second dose of antigen in mice.

Method

The materials and methods for mouse pinnal anaphylaxis were previously described except that all samples of Beta glucuronidase had been passed through a column of polyacrilamide gel (Bio-Gel P-6 Bio Rad Lab) to remove the sorbitol which was added to the enzyme as a stabilizer during commercial freeze-drying. For this purpose the buffer previously described was employed without phosphate. The protein void from the column was saved. There was no appreciable loss of total Beta glucuronidase activity. Sterility was maintained by passage through Millipore filters. Gelatin was obtained from Armour Pharmaceutical Co. Ltd. and from E. Merck A.G. Solutions were stored at -20°C before use.

The "in vitro" assay of Beta glucuronidase activity using phenolphthalein glucuronide (Sigma Chemical Co.) was carried out by the method described in Sigma technical Bulletin No. 105 (1958). However, the experiments were duplicated in the buffer used for the animal work which has a pH of 5-6. Further experiments assessed the effect of glucose and N-acetyl glucosamine on the plasma inhibitor of Beta glucuronidase,

using the same buffer with 6.5 percent defibrinated human serum.

Results

In an earlier study, it was noticed that batches of Beta glucuronidase which had aged lost their hyposensitizing activity and often had the opposite effect. Table 1 shows the effect of such a sample of enzyme, freeze-dried and stored at -80°C for 18 months, on the anamnestic response of mice.

The addition of this aged enzyme to the second dose of antigen results in a repeatable, but not significant, increase in anaphylactic sensitivity. When glucose 0.1 gamma g per mouse is added to the treatment mixture of antigen and enzyme shortly before injection, the ability of the enzyme to block the anamnestic response is revealed.

If concentrations of glucose greater and less than 0.1 gamma g are added to the enzyme and antigen mixture, they induce an increase in anaphylactic sensitivity, but if glucose and enzyme are allowed to interact "in vitro" at room temperature for 1 hour before the mixture is injected, the critical dose of glucose for blockade of the anamnestic response is reduced tenfold to 0.01 gamma g/mouse.

This ability to modify the effect of Beta glucuronidase is not peculiar to glucose; the action is shared by many other sugars. Figure 1 shows the striking similarity of the dose-response curves obtai-

ned with varying concentrations of glucose, glucosamine, N-acetyl glucosamine, N-acetyl galactosamine and N-acetyl neuraminic acid. The results are expressed as percent change of the effect of antigen and enzyme without added sugar. The five dose-response curves shown are taken from consecutive experiments.

In this series, the sample of Beta glucuronidase used had a slight hypersensitizing effect when not modified by sugar. Table II shows the effect of adding gelatin 1.0 ng per mouse to the treatment mixture shortly before injection. In this experiment the Beta glucuronidase had not been stored for a long period, and possessed the ability to block the anamnestic response without the addition of glucose. Gelatin reverses the action of the enzyme so that it induces hypersensitization. This result was reproduced in five consecutive experiments. The gelatin from Merck had a consistent effect over a wide dose range. The gelatin from Armour also blocked the desired effect of the enzyme at 1.0 ng per mouse, but at higher concentrations its dose-response curve followed a pattern which resembled those of the sugars in figure 1.

Table III shows that glucose can restore the ability to block the anamnestic response to a mixture of enzyme and gelatin. In further experiments glucose was shown to antagonize the effect of both samples of gelatin.

Experiments "in vitro" have been carried out to assess the effect of glucose, N-

Table 1. Pinnal anaphylaxis in groups of 36 mice. Activation by glucose 0.1 μg of the ability of aged β -glucuronidase 10 U, to block the anamnestic response to a subcutaneous dose of 1 μg horse serum given 3 weeks after initial sensitization and 8 days before challenge

Treatment	Mean area blued mm
Saline	29
Horse serum	36
Horse serum + β -glucuronidase	40 ^a
Horse serum + β -glucuronidase + glucose	30 ^a

^a Significance: $p = < 0.02$.

Table 2. Pinnal anaphylaxis in groups of 21 mice. Effect of gelatin 1.0 ng on the ability of β -glucuronidase 10 U to block the anamnestic response to a subcutaneous dose of 1 μg horse serum given 3 weeks after initial sensitization and 8 days before challenge

Treatment	Mean area blued mm
Saline	46
Horse serum	56
Horse serum + β -glucuronidase	47 ^a
Horse serum + β -glucuronidase + gelatin	62 ^a

^a Significance: $p = < 0.02$.

Table 3. Pinnal anaphylaxis in groups of 8 mice. Effect of glucose 1.0 μ g and gelatin 1.0 ng on the ability of β -glucuronidase 10 U to block the anamnestic responses to a subcutaneous dose of 1 μ g horse serum given 3 weeks after initial sensitization and 8 days before challenge

Treatment	Mean area blued mm
Saline	40
Horse serum	51
Horse serum + β -glucuronidase	41
Horse serum + β -glucuronidase + gelatin	57 ^a
Horse serum + β -glucuronidase + gelatin + glucose	40 ^a

^a Significanos: $p = < 0.06$.

acetyl glucosamine and gelatin on the activity of the enzyme against phenolphthalein glucuronide. At concentrations of 1 percent, glucose and N-acetyl glucosamine inhibit the enzyme's activity by 15 percent and 36 percent respectively.

Greater dilutions have little effect. Defibrinated human serum 6.5 percent reduces the enzyme's activity to 20 percent of the control, but N-acetyl glucosamine 1 percent restores the activity to 33 percent of control. Weak antagonism of the plasma inhibitor of Beta glucuronidase is seen with concentrations of N-acetyl glucosamine down to 0.01 percent. Glucose does not behave in this way. Gelatin has no influence on the activity of the Beta glucuronidase used in the present experiments.

Discussion

These experiments identify an obstacle which has prevented further progress in the study of enzyme-potentiated hyposensitization for a number of years. Different samples of Beta glucuronidase have widely differing hyposensitizing effects. Some batches of "Patella Vulgata" enzyme have no hyposensitizing activity even when fresh, and all batches will slowly lose their ability in this direction on storage, although freeze-dried and kept at -80°C . Nevertheless, the activities of different samples against




Fig. 1. Mouse pinnal anaphylaxis: groups of seven mice sensitized with 250 μ g horse serum. Treated after three weeks with 1 μ g horse serum + 10 U β -glucuronidase + various doses of sugars and challenged eight days later. Results expressed as percentage of result after treatment with antigen + enzymes.

phenolphthalein glucuronide do not differ greatly, and their degrees of purity do not vary sufficiently to account for their great differences in hyposensitizing ability.

The two most likely explanations of these facts are either that a contaminant of the Beta glucuronidase is really responsible for its hyposensitizing effect, or that contaminants, perhaps other enzymes, interfere with the desired activity. The latter explanation commended itself early in this work when Robinson and Stirling examined samples of ovine testicular hyaluronidase and found that batches which possessed hyposensitizing activity contained more Beta glucuronidase than N-acetyl glucosamine. In batches with no hyposensitizing activity the proportions were reversed.

The first possibility has also been considered since Mowbray and his coworkers have reported that polyribonuclease is capable of inducing immunological tolerance, and the Beta glucuronidase used in the present work was contaminated with ribonuclease. There are, however, a number of differences between the beha-

viour of Mowbray's enzyme preparation and the Beta glucuronidase used in the present experiments: First, in Mowbray's work, tolerance induction by polyribonuclease could not be achieved unless it was given some hours before or after the antigen. Administration by a protocol similar to that used for Beta glucuronidase was ineffective. Second, the optimal dose of purified polyribonuclease was 200 gamma g per mouse. This compares with 5 gamma g per mouse for impure Beta glucuronidase. Third, very small doses of sugars would not be expected to influence the behaviour of polyribonuclease, but suggest that a carbohydrase or a carbohydrate receptor are more likely to be involved.

Newsome has found that the tolerance-inducing properties of polyribonuclease preparations were due to a product of contaminating bacteria and not to the enzyme itself. The Beta glucuronidase used in the present work was purchased already freeze-dried, and was sterile at all stages thereafter.

Original contamination of the molluscan digestive glands is a possibility which has not been excluded. Nevertheless, the difference in behaviour between Beta glucuronidase and the agent investigated by Mowbray strongly suggest that they are not identical.

The work reported here shows that the ability of Beta glucuronidase from "Patella vulgata" to block the anamnestic response in mice is activated by the presence of extremely small quantities of sugars. In earlier work, Beta glucuronidase was used with sorbitol 20 percent by weight as stabilizer. It seems possible that the variability of results at this time may have been partly due to contamination of the commercial sorbitol with glucose or other sugars. At the above concentration, pure sorbitol dose not cause Beta glucuronidase to block the anamnestic response in mice, and the ability of added glucose to elicit this activity from aged Beta glucuronidase was first noted in four consecutive mouse experiments using enzyme which contained sorbitol.

Figure 1 shows that reversal of the effect of the enzyme and antigen mixture can be brought about by varying the dose of sugar employed. The unusual dose-response relationship will be considered more fully in the next publication in this series.

"In vitro", neither glucose nor N-acetyl glucosamine altered the enzyme's hydrolysis of phenolphthalein glucuronide except in concentration of 1 percent, when some suppression of enzyme activity was noted. This contrasts with the 10,000-fold lower dose of either required to potentiate the enzyme's ability to block the anamnestic response in the mouse.

The inhibitory effect of plasma on the enzyme's activity "in vitro" was antagonized to a small degree by N-acetyl glucosamine, again only at high concentrations. Glucose did not possess this action. Since glucose and N-acetyl glucosamine have similar effects on the immunological behaviour of the enzyme, it is inferred that for this role, any action in the plasma inhibitor of Beta glucuronidase is immaterial.

Gelatin is an activator of purified Beta glucuronidase "in vitro". At the low degree of purity of the enzyme used in the present experiments, gelatin had no effect on "in vitro" activity against phenolphthalein glucuronide. "In vivo", gelatin's ability to alter the hyposensitizing effect of the enzyme was studied since it seemed likely that other denatured proteins might act in the same way. In common with many proteins, gelatin is known to contain many hexose groups. It is suggested that the difference in behaviour of the two samples which were tested was due to some difference in the distribution of the hexose groups on the surface of the molecules.

Molluscan Beta glucuronidase from *Helix pomatia* has been prepared with an activity of 120,000 U/mg, by which standard the preparation used in the present work contained about 98 percent of extraneous protein. The precipitation procedure used in the present work of the enzyme, and ageing after

freeze-drying might both produce variable amounts of altered contaminants capable of exerting a gelatin-like effect. Since gelatin has this action at such low concentrations, it is clear that a very high degree of purification of the enzyme would be necessary to obviate the possibility. That the enzyme's loss of hyposensitizing activity on storage is in reality due to an alteration of a contaminant is suggested by the ability of glucose to restore the anti-anamnestic effect

to aged enzyme as well as to reverse the action of gelatin.

Acknowledgements

Part of this work was supported by grants from the Miriam Marks Charitable Trust and from the Asthma Research Council to whom I am grateful. Again I thank Miss Mary Fitzgerald for her indispensable technical help.